



eVade™

Crabgrass Preventer with EVADE™ .375% Plus

ACTIVE INGREDIENT:
Proflumicafene (1,3,5-tri-*n*-propyl-2,4-dinitro-6-
(trifluoromethyl)-*m*-phenylene diamine) 0.375%
INERT INGREDIENTS: 99.625%
TOTAL: 100.000%

For selective preemergence control of grass and broadleaf weeds in:

- Established turf grasses (excluding golf course putting greens) and lawns
- Landscape ornamentals (including established perennials & wildflower plantings)
- Covers up to 16,390 square feet

KEEP OUT OF REACH OF CHILDREN

CAUTION

See the following section, Precautionary Statements,
Hazards to Humans and Domestic Animals, for additional information.

EPA REG. NO. 34704-776

EPA EST. NO. 84156-CAN-2 (Lot begins 2) EPA EST. NO. 41124-OH-1 (Lot begins 0)
EPA EST. NO. 41124-WI-1 (Lot begins W) EPA EST. NO. 49991-TX-1 (Lot begins TX-1)
EPA EST. NO. 57131-WI-1 (Lot begins EC) EPA EST. NO. 49991-TX-2 (Lot begins TX-2)
EPA EST. NO. 75521-PA-12 (Lot begins FS) EPA EST. NO. 82757-MA-2 (Lot begins H)
EPA EST. NO. 82757-FL-1 (Lot begins S) EPA EST. NO. 82757-OH-1 (Lot begins M)
NET WEIGHT 50 LBS. (22.68 KG)

III

EXP 01/10

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact while handling the material may cause allergic reactions in some individuals.

FIRST AID

If in eyes:	• Hold eye open and rinse slowly and gently with water for 15–20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	• Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15–20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	• Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. • Call a poison control center or doctor for further treatment advice.
If swallowed:	• Call a poison control center or doctor immediately for treatment advice. • Have a person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.

FOR A MEDICAL EMERGENCY INVOLVING THIS PRODUCT CALL: 1-866-944-8565.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

ENVIRONMENTAL HAZARDS

This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

If material is spilled: Contain and sweep up material of spill for disposal or use per label instructions. Wash cleanup equipment to prevent contamination elsewhere. Excess quantities above normal application rates could result in damage to plants subject to spill. Water in remainder, disc under, cover with layer of soil, or leave exposed to sunlight, as deemed appropriate to the situation.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

STORAGE: Store in original container away from other fertilizer, feed, or food stuffs and separated from other pesticides.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Nonrefillable container. Do not reuse or refill this container. Completely empty bag into application equipment by shaking and tapping sides and bottom to loosen clinging particles. Offer for recycling if available or dispose of empty bag in a sanitary landfill or by incineration. Do not burn, unless allowed by state and local ordinances.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This product is a selective preemergence herbicide that provides residual control of many grass and broadleaf weeds in:

- established turf grasses (excluding golf course putting greens) and lawns
- landscape ornamentals
- established perennials & wildflower plantings

This product controls susceptible weeds by inhibiting weed seeds, germination and root development. Most effective weed control will be obtained when this product is activated by at least 0.5 inches of rainfall, or irrigation, or with shallow (1 to 2 inches) incorporation, prior to weed seed germination and within 14 days following application.

Not for use on plants being grown (1) for sale or other commercial use, (2) for commercial seed production, or (3) for research purposes. For use on plants intended for aesthetic purposes or climatic modification and being grown in ornamental gardens or parks, or on golf courses, or lawns and grounds.

Do not graze or feed livestock forage cut from areas treated with this product.

Do not apply this product through any type of irrigation system.

Do not apply aerially.

Do not apply to golf course putting greens.

Apply uniformly with suitable, calibrated application equipment.

APPLICATION DIRECTIONS

GUARANTEED ANALYSIS:

Total Nitrogen (N) 13.00%
13.00% Urea Nitrogen*
Soluble Potash (K₂O) 4.00%
DERIVED FROM: Polymer Coated Sulfur Coated Urea, Urea, Murate of Potash.
*2.42% Slowly available Nitrogen from Polymer Coated Sulfur Coated Urea.

WARNING: THIS PRODUCT CONTAINS CHEMICALS KNOWN TO THE STATE OF CALIFORNIA TO CAUSE CANCER OR BIRTH DEFECTS OR OTHER REPRODUCTIVE HARM

Information regarding the contents and level of metals in this product is available on the Internet at <http://www.regulatory-info-lpi.com>

Product No. 8112380

GENERAL APPLICATION GUIDELINES:

Rate	lbs. Product per 1,000 sq ft	lbs. Nitrogen per 1,000 sq ft	lbs. A.I. per acre	
Low	3.05	0.40	0.50	
Medium	4.60	0.60	0.75	
High	6.14	0.80	1.00	
Spreader Setting	Bigfoot	Earthway	Lesco Rotary	AcurPro 2000
Low	L	16	J	N
Medium	N	17	K	O
High	P	18	L	P

These settings are approximate for the application rate shown per 1,000 square feet. You may need to adjust the spreader setting depending on walking speed, spreader condition and product.

Contains XCU Polymer Coated Sulfur Coated Urea, a trademark of Agrinum Advanced Technologies.



Slow Release Fertilizer

ESTABLISHED TURF


This product is a selective preemergence herbicide that, when properly applied, will control certain grass and broadleaf weeds in established turf grasses and lawns. The maximum amount of this product that may be applied per year is given for each turf grass species in the Medium Annual Rates section of this label. Most



DISTRIBUTED BY
LOVELAND PRODUCTS, INC.
P.O. BOX 1286, GREELEY, COLORADO 80632-1286

Signature Crabgrass Preventer with EVADE .375% Plus
EPA REG. NO. 34704-776

CRAAGGRASS SEED GERMINATION DATES



crabgrass germination information for your area.

Use Precautions

The following precautions apply to the use of this product in turf grasses and lawns: Application of this product may thin emerged annual bluegrass and newly overseeded grasses.

in the second inch of soil, not thatch plus soil

following year.

or annual bluegrass (*Poa annua*) are desirable species

Rates of Application

uct will not control established weeds.

provided by this product is related to the rate applied

Maximum Annual Rates

maximum yearly rate as given in the following table:

Maximum Application Rate (Calendar Year of Crabgrass Preventer with ENADE 375% Plus by Turf Grass Species) ¹		
Turf Species:	lbs product/A	lbs a.i./A
Bermudagrass ^{2b}	400	1.5
Bahia grass		
Centipedegrass		
Seashore Paspalum		
St. Augustinegrass		
Tall Fescue (including turf-type)		
Zoysia		
Buffalograss	267	1.0
Kentucky Bluegrass		
Perennial Ryegrass		
Cureeping Red Fescue	200	0.75

LANDSCAPE ORNAMENTALS

This product may be applied for residual preemergence weed control in ornamentals.

Use Rates

does not exceed the maximum annual application rate of 400 lbs/A/yr

Application Timing and Information

in ornamentals will be obtained when this product is activated by at least 0.5 inches of rainfall or irrigation.

[illegible]

use of an appropriate postemergence herbicide

Established Landscape Ornamentals

the soil

Newly Transplanted Landscape Ornamentals

ing or after buds/grafts have taken to avoid any inhibition of the tissue union

Tolerant Ornamental Species

sitive to this product

Not for use on container grown plants.

CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

or agent of LOVELAND PRODUCTS, INC. or the seller is authorized to vary in any way

or user of this product assumes all such inherent risks

PURPOSE, OR ELIGIBILITY OF THIS PRODUCT FOR ANY PARTICULAR TRADE USAGE

UCTS, INC., ATTENTION: LAW DEPARTMENT, P.O. BOX 1286, GREELEY, CO 80632-1286

SEQUENTIAL, SPECIAL, OR INDIRECT DAMAGES, OR DAMAGES IN THE NATURE OF A PENALTY.

EVAADE is a trademark of Loveland Products, Inc.